

REFERENCIAS BIBLIOGRÁFICAS

1. Chan S, McCowen KC, Blackburn GL. Nutrition management in the ICU. *Chest* 1999;115(5 Suppl):S145-S148.
2. Salazar O. Estar críticamente enfermo significa no ser capaz y no poder decidir. *Index Enferm* 2008;17:5.
3. Hickman RL Jr, Douglas SL. Impact of chronic critical illness on the psychological outcomes of family members. *AACN Adv Crit Care* 2010;21:80-91.
4. Valero ZM, Alvarez NR, García RP, Sánchez GR, Moreno VJ, León SM. Should hydration and artificial nutrition be accepted as a palliative care? *Nutrición Hospitalaria [España]* 2006;21:680-5.
5. DeLegge MH, McClave SA, DiSario JA, Baskin WN, Brown RD, Fang JC; et al; for the ASGE Task Force on Enteral Nutrition. Ethical and medicolegal aspects of PEG-tube placement and provision of artificial nutritional therapy. *Gastrointest Endosc* 2005;62:952-9.
6. Van der Riet P, Brooks D, Ashby M. Nutrition and hydration at the end of life: Pilot study of a palliative care experience. *J Law Med* 2006;14:182-98.
7. Plank LD, Hill GL. Sequential metabolic changes following induction of systemic inflammatory response in patients with severe sepsis or major blunt trauma. *World J Surg* 2000;24:630-8.
8. Streat SJ, Beddoe AH, Hill GL. Aggressive nutritional support does not prevent protein loss despite fat gain in septic intensive care patients. *J Trauma* 1987;27:262-6.
9. Montejo González JC, Culebras Fernández JM, García de Lorenzo y Mateos A. Recommendations for the nutritional assessment of critically ill patients. *Rev Med Chil* 2006;134:1049-56.
10. Bistrian BR. A simple technique to estimate severity of stress. *Surg Gynecol Obstet* 1979; 148:675-8.
11. Bruyninckx F, Schetz M, Vlasselaers D, Ferdinand P, Lauwers P, Bouillon R. Intensive insulin therapy in the critically ill patients. *New Engl J Med* 2001;345:1353-67.
12. van den Berghe G, Wilmer A, Hermans G, Meersseman W, Wouters PJ, Milants I; et al. Intensive insulin therapy in the medical ICU. *New Engl J Med* 2006;354:449-61.
13. Krinsley JS. Effect of an intensive glucose management protocol on the mortality of critically ill adult patients. *Mayo Clin Proc* 2004;79:992-1000.
14. Montecalvo MA, Steger KA, Farber HW, Smith BF, Dennis RC, Fitzpatrick GF; et al; for the The Critical Care Research Team. Nutritional outcome and pneumonia in critical care patients randomized to gastric versus jejunal tube feedings. *Crit Care Med* 1992;20:1377-87.
15. Kompan L, Kremžvar B, Gadžvijev E, Prošek M. Effects of early enteral nutrition on intestinal permeability and the development of multiple organ failure after multiple injury. *Intensive Care Med* 1999;25:157-61.
16. Heyland DK, Dhaliwal R, Drover JW, Gramlich L, Dodek P. Canadian clinical practice guidelines for nutrition support in mechanically ventilated, critically ill adult patients. *JPEN J Parenter Enteral Nutr* 2003;27:355-373. Actualización disponible *on-line* en: <http://www.criticalcarenutrition.com>. Fecha de última actualización: 13 de Diciembre del 2015.

17. McClave SA, Martindale RG, Vanek VW, McCarthy M, Roberts P, Taylor B; *et al*; for the American College of Critical Care Medicine. Guidelines for the provision and assessment of nutrition support therapy in the adult critically ill patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (ASPEN). *JPEN J Parenter Enteral Nutr* 2009;33:277-316.
18. Martín Delgado M, Saura R, Cabré Pericas L, Ruiz Moreno J, Blanch L, Blanco J; for the Working Group of the Sociedad Española de Medicina Intensiva, Crítica y Unidades Coronarias (SEMICYUC). Quality indicators in critically ill patients. *Critical Care* 2006; 10(Suppl 1):P395.
19. Kreymann KG, Berger MM, Deutz NE, Hiesmayr M, Jollet P, Kazandjiev G; *et al*. ESPEN guidelines on enteral nutrition: Intensive care. *Clinical Nutrition* 2006;25:210-23.
20. Singer P, Berger MM, van den Berghe G, Biolo G, Calder P, Forbes A; *et al*. ESPEN guidelines on parenteral nutrition: Intensive care. *Clinical Nutrition* 2009;28:387-400.
21. ADA American Dietetic Association. Critical illness evidence-based nutrition practice guideline. Washington DC: 2006. Disponible en: <http://www.adaevidencelibrary.com/topic.cmf>. Fecha de última visita: 14 de Diciembre del 2015.
22. Doig GS, Simpson F; for the Australian and New Zealand Intensive Care Society Clinical Trials Group. Evidence-based guidelines for nutritional support of the critically ill: Results of a bi-national guideline development conference. ANZICS Australian and New Zealand Intensive Care Society. Carlton (Australia): 2005. Disponible en: <http://EvidenceBased.net>. Ficha de última visita: 23 de Diciembre del 215.
23. Mesejo A , Vaquerizo Alonso C , Acosta Escribano J , Ortiz Leyba C, Montejo González JC; *et al*; for the Spanish Society of Intensive Care Medicine and Coronary Units and the Spanish Society of Parenteral and Enteral Nutrition (SEMICYUC-SENPE). Guidelines for specialized nutritional and metabolic support in the critically-ill patient. Update. Consensus of the Spanish Society of Intensive Care Medicine and Coronary Units and the Spanish Society of Parenteral and Enteral Nutrition (SEMICYUC-SENPE): Introduction and methodology. *Medicina Intensiva [España]* 2011;35(Suppl 1):1-6.
24. Terrace L. Development and validation of an international appraisal instrument for assessing the quality of clinical practice guidelines: The AGREE project. *Qual Saf Health Care* 2003; 12:18-23.
25. MinSal Ministerio de Salud de la Nación. Estandarización de procesos asistenciales. Calidad de gestión clínica. República Argentina. Buenos Aires: 2007.
26. Acosta Escribano J, Gómez-Tello V, Ruiz Santana S. Nutritional assessment of the severely ill patient. *Nutrición Hospitalaria [España]* 2005;20(Suppl 2):S5-S8.
27. Mesejo A. Recomendaciones para el soporte nutricional del paciente crítico. *Medicina Intensiva [España]* 2011;35(Supl 1):S85-S91.
28. Goiburu ME, Goiburu MM, Bianco H, Díaz JR, Alderete F, Palacios MC; *et al*. The impact of malnutrition on morbidity, mortality and length of hospital stay in trauma patients. *Nutrición Hospitalaria [España]* 2006;21:604-10.
29. Sungurtekin H, Sungurtekin U, Oner O, Okke D. Nutrition assessment in critically ill patients. *Nutr Clin Pract* 2008;23:635-41.
30. Fontes D, de Vasconcelos Generoso S, Correia MITD. Subjective global assessment: A reliable nutritional assessment tool to predict outcomes in critically ill patients. *Clin Nutr* 2014;33:291-5.

31. Heyland DK, Dhaliwal R, Jiang X, Day A. G. Identifying critically ill patients who benefit the most from nutrition therapy: The development and initial validation of a novel risk assessment tool. *Crit Care* 2011;15:R268.
32. Choban P, Dickerson R, Malone A, Worthington P, Compher C. ASPEN clinical guidelines: Nutrition support of hospitalized adult patients with obesity. *JPEN J Parenter Enteral Nutr* 2013;37:714-44.
33. Rahman A, Hasan RM, Agarwala R, Martin C, Day AG, Heyland DK. Identifying critically-ill patients who will benefit most from nutritional therapy: Further validation of the "modified NUTRIC" nutritional risk assessment tool. *Clin Nutr* 2016;35:158-62.
34. Kang W, Kudsk KA. Is there evidence that the gut contributes to mucosal immunity in humans? *JPEN J Parenter Enteral Nutr* 2007;31:246-58.
35. Windsor ACJ, Kanwar S, Li AGK, Barnes E, Guthrie JA, Spark JI; *et al.* Compared with parenteral nutrition, enteral feeding attenuates the acute phase response and improves disease severity in acute pancreatitis. *Gut* 1998;42:431-5.
36. Ammori B, Leeder P, King R, Barclay R, Martin I, Larvin M, McMahon M. Early increase in intestinal permeability in patients with severe acute pancreatitis: Correlation with endotoxemia, organ failure, and mortality. *J Gastrointest Surg* 1999;3:10-8.
37. Nguyen NQ, Fraser RJ, Bryant LK, Burgstad C, Chapman MJ, Bellon M; *et al.* The impact of delaying enteral feeding on gastric emptying, plasma cholecystokinin, and peptide YY concentrations in critically ill patients. *Crit Care Med* 2008;36:1469-74.
38. Chiarelli A, Enzi G, Casadei A, Baggio B, Valerio A, Mazzoleni F. Very early nutrition supplementation in burned patients. *Am J Clin Nutr* 1990;51:1035-9.
39. Grahm TW, Zadrozny DB, Harrington T. The benefits of early jejunal hyperalimentation in the head-injured patient. *Neurosurgery* 1989;25:729-35.
40. Taylor S, Fettes SB, Jewkes C, Nelson R. Prospective, randomized, controlled trial to determine the effect of early enhanced enteral nutrition on clinical outcome in mechanically ventilated patients suffering head injury. *Crit Care Med* 1999;27:2525-2531.
41. Singh G, Ram RP, Khanna SK. Early postoperative enteral feeding in patients with nontraumatic intestinal perforation and peritonitis. *J Am Coll Surg* 1998;187:142-6.
42. Moore EE, Jones TN. Benefits of immediate jejunostomy feeding after major abdominal trauma- A prospective, randomized study. *J Trauma* 1986;26:874-81.
43. Khalid I, Doshi P, DiGiovine B. Early enteral nutrition and outcomes of critically ill patients treated with vasopressors and mechanical ventilation. *Am J Crit Care* 2010;19:261-8.
44. Eyer SD, Micon LT, Konstantinides FN, Edlund DA, Rooney KA, Luxenberg MG, Cerra FB. Early enteral feeding does not attenuate metabolic response after blunt trauma. *J Trauma* 1993;34:639-43.
45. Ibrahim EH, Mehringer L, Prentice D, Sherman G, Schaiff R, Fraser V, Kollef MH. Early versus late enteral feeding of mechanically ventilated patients: Results of a clinical trial. *JPEN J Parenter Enteral Nutr* 2002;26:174-81.
46. Marik PE, Zaloga GP. Early enteral nutrition in acutely ill patients: A systematic review. *Crit Care Med* 2001;29:2264-70.
47. Doig GS, Heighes PT, Simpson F, Sweetman EA, Davies AR. Early enteral nutrition, provided within 24 h of injury or intensive care unit admission, significantly reduces mortality in critically ill patients: A meta-analysis of randomised controlled trials. *Intensive Care Med* 2009; 35:2018-27.

48. Chourdakis M, Kraus MM, Tzellos T, Sardeli C, Peftoulidou M, Vassilakos D, Kouvelas D. Effect of early compared with delayed enteral nutrition on endocrine function in patients with traumatic brain injury: An open-labeled randomized trial. JPEN J Parenter Enteral Nutr 2012;36:108-16.
49. Zaloga GP, Roberts PR, Marik P. Feeding the hemodynamically unstable patient: A critical evaluation of the evidence. Nutr Clin Pract 2003;18:285-93.
50. Mutlu GM, Mutlu EA, Factor P. Prevention and treatment of gastrointestinal complications in patients on mechanical ventilation. Am J Respir Med 2003;2:395-411.
51. Dvorak MF, Noonan VK, Bélanger, L, Bruun B, Wing PC, Boyd MC, Fisher C. Early versus late enteral feeding in patients with acute cervical spinal cord injury: A pilot study. Spine [Philadelphia] 2004;29:E175-E180.
52. Casaer MP, van den Berghe G. Nutrition in the acute phase of critical illness. New Engl J Med 2014;370:1227-1236.
53. Braunschweig CL, Levy P, Sheean PM, Wang X. Enteral compared with parenteral nutrition: A meta-analysis. Am J Clin Nutr 2001;74:534-42.
54. Peter JV, Moran JL, Phillips-Hughes J. A metaanalysis of treatment outcomes of early enteral versus early parenteral nutrition in hospitalized patients. Crit Care Med 2005;33:213-20.
55. Moore FA, Feliciano DV, Andrassy RJ, McArdle AH, Booth FW, Morgenstein-Wagner TB; *et al.* Early enteral feeding, compared with parenteral, reduces postoperative septic complications. The results of a meta-analysis. Ann Surg 1992;216:172-83.
56. Simpson F, Doig GS. Parenteral vs. enteral nutrition in the critically ill patient: A meta-analysis of trials using the intention to treat principle. Intensive Care Med 2005;31:12-23.
57. Lipman TO. Grains or veins: Is enteral nutrition really better than parenteral nutrition? A look at the evidence. JPEN J Parenter Enteral Nutr 1998;22:167-82.
58. Gramlich L, Kichian K, Pinilla J, Rodych NJ, Dhaliwal R, Heyland DK. Does enteral nutrition compared to parenteral nutrition result in better outcomes in critically ill adult patients? A systematic review of the literature. Nutrition 2004;20:843-8.
59. Casaer MP, Mesotten D, Hermans G, Wouters PJ, Schetz M, Meyfroidt G; *et al.* Early versus late parenteral nutrition in critically ill adults. N Engl J Med 2011;365:506-17.
60. Dhaliwal R, Jurewitsch B, Harrietha D, Heyland DK. Combination enteral and parenteral nutrition in critically ill patients: Harmful or beneficial? A systematic review of the evidence. Intensive Care Med 2004;30:1666-71.
61. Heyland DK, MacDonald S, Keefe L, Drover JW. Total parenteral nutrition in the critically ill patient: A metaanalysis. JAMA 1998;280:2013-9.
62. Krishnan JA, Parce PB, Martinez A, Diette GB, Brower RG. Caloric intake in medical ICU patients: consistency of care with guidelines and relationship to clinical outcomes. Chest 2003;124:297-305.
63. Villet S, Chiolero RL, Bollmann MD, Revelly JP, Cayeux MC, Delarue J, Berger MM. Negative impact of hypocaloric feeding and energy balance on clinical outcome in ICU patients. Clin Nutr 2005;24:502-9.
64. Harris JA, Benedict FG. A biometric study of human basal metabolism. PNAS Proc Nat Acad Sci 1918;4:370-3.
65. Schofield WN, Schofield C, James WPT. Basal metabolic rate: Review and prediction, together with an annotated bibliography of source material. Human Nutrition Clinical Nutrition 1985;39(Suppl 1):42-91.

66. Ireton-Jones CS, Turner Jr WW, Liepa GU, Baxter CR. Equations for the estimation of energy expenditures in patients with burns with special reference to ventilatory status. *J Burn Care Research* 1992;13:330-3.
67. Boullata J, Williams J, Cottrell F, Hudson L, Compher C. Accurate determination of energy needs in hospitalized patients. *J Am Diet Assoc* 2007;107:393-401.
68. Flancbaum L, Choban PS, Sambucco S, Verducci J, Burge JC. Comparison of indirect calorimetry, the Fick method, and prediction equations in estimating the energy requirements of critically ill patients. *Am J ClinNutr* 1999;69:461-6.
69. Saffle JR, Larson CM, Sullivan J. A randomized trial of indirect calorimetry based feedings in thermal injury. *J Trauma* 1990;30:776-82.
70. Haugen HA, Chan LN, Li F. Indirect calorimetry: A practical guide for clinicians. *Nutr Clin Pract* 2007;22:377-88.
71. Swinamer DL, Phang PT, Jones RL, Grace M, King EG. Twenty-four hour energy expenditure in critically ill patients. *Crit Care Med* 1987;15:637-43.
72. Ireton-Jones C. Adjusted body weight, con: Why adjust body weight in energy expenditure calculations? *Nutr Clin Pract* 2005;20:474-9.
73. Arabi YM, Haddad SH, Tamim HM, Rishu AH, Sakkijha MH, Kahoul SH, Britts RJ. Near-target caloric intake in critically ill medical-surgical patients is associated with adverse outcomes. *JPEN J Parenter Enteral Nutr* 2010;34:280-8.
74. Singer P, Anbar R, Cohen J, Shapiro H, Shalita-Chesner M, Lev S. The tight calorie control study (TICACOS): A prospective, randomized, controlled pilot study of nutritional support in critically ill patients. *Intensive Care Med* 2011;37:601-9.
75. Frankenfield DC, Coleman A, Alam S, Cooney RN. Analysis of estimation methods for resting metabolic rate in critically ill adults. *JPEN J Parenter Enteral Nutr* 2009;33:27-36.
76. Frankenfield D. Validation of an equation for resting metabolic rate in older obese, critically ill patients. *JPEN J Parenter Enteral Nutr* 2011;35:264-9.
77. Desachy A, Clavel M, Vuagnat A, Normand S, Gissot V, François B. Initial efficacy and tolerability of early enteral nutrition with immediate or gradual introduction in intubated patients. *Intensive Care Med* 2008;34:1054-9.
78. Martin CM, Doig GS, Heyland DK, Morrison T, Sibbald WJ; for the Southwestern Ontario Critical Care Research Network. Multicentre, cluster-randomized clinical trial of algorithms for critical-care enteral and parenteral therapy (ACCEPT). *CMAJ* 2004;170:197-204.
79. Doig GS, Simpson F, Finfer S, Delaney A, Davies AR, Mitchell I; *et al*; for the Nutrition Guidelines Investigators of the ANZICS Clinical Trials Group. Effect of evidence-based feeding guidelines on mortality of critically ill adults: A cluster randomized controlled trial. *JAMA* 2008;300:2731-41.
80. Rice TW, Mogan S, Hays MA, Bernard GR, Jensen GL, Wheeler AP. Randomized trial of initial trophic versus full-energy enteral nutrition in mechanically ventilated patients with acute respiratory failure. *Crit Care Med* 2011;39:967-74.
81. Rice TW, Wheeler AP, Thompson BT, Steingrub J, Hite RD, Moss M; *et al*. Initial trophic vs. full enteral feeding in patients with acute lung injury: The EDEN randomized trial. *JAMA* 2012;307:795-803.
82. Guía de Práctica Clínica de Soporte Nutricional Enteral y Parenteral en Pacientes Hospitalizados y Domiciliarios. Programa Nacional de Garantía de la Calidad de la Atención Médica. MINSAL Ministerio de Salud de la Nación. República Argentina. Buenos Aires: 2007. pp. 47-55.

83. Dickerson RN, Boschert KJ, Kudsk KA, Brown RO. Hypocaloric enteral tube feeding in critically ill obese patients. *Nutrition* 2002;18:241-6.
84. Alberda C, Gramlich L, Jones N, Jeejeebhoy K, Day AG, Dhaliwal R, Heyland DK. The relationship between nutritional intake and clinical outcomes in critically ill patients: Results of an international multicenter observational study. *Intensive Care Med* 2009;35:1728-37.
85. Liu KJ, Cho MJ, Atten MJ, Panizales E. Hypocaloric parenteral nutrition support in elderly obese patients. *Am Surg* 2000;66:394-9.
86. Dickerson RN, Medling TL, Smith AC, Maish GO, Croce MA, Minard G, Brown RO. Hypocaloric, high-protein nutrition therapy in older vs. younger critically ill patients with obesity. *JPEN J Parenter Enteral Nutr* 2013;37:342-51.
87. Mentec H, Dupont H, Bocchetti M, Cani P, Ponche F, Bleichner G. Upper digestive intolerance during enteral nutrition in critically ill patients: Frequency, risk factors, and complications. *Crit Care Med* 2001;29:1955-61.
88. Ritz MA, Fraser R, Tam W, Dent J. Impacts and patterns of disturbed gastrointestinal function in critically ill patients. *Am J Gastroenterol* 2000;95:3044-52.
89. Scolapio JS. Methods for decreasing risk of aspiration pneumonia in critically ill patients. *JPEN J Parenter Enteral Nutr* 2002;26(6 Suppl):S58-S61.
90. Deane A, Chapman M, Fraser R, Bryant L, Burgstad C, Nguyen Q. Mechanisms underlying feed intolerance in the critically ill: Implications for treatment. *World J Gastroenterol* 2007; 13:3909-17.
91. Dive A, Moulart M, Jonard P, Jamart J, Mahieu P. Gastroduodenal motility in mechanically ventilated critically ill patients: A manometric study. *Crit Care Med* 1994;22:441-7.
92. Fruhwald S, Kainz J. Effect of ICU interventions on gastrointestinal motility. *Curr Opin Crit Care* 2010;16:159-64.
93. McClave SA, DeMeo MT, DeLegge MH, DiSario JA, Heyland DK, Maloney JP; *et al.* North American Summit on Aspiration in the Critically Ill Patient: Consensus statement. *JPEN J Parenter Enteral Nutr* 2002;26(6 Suppl):S80-S5.
94. Moreira TV, McQuiggan M. Methods for the assessment of gastric emptying in critically ill, enterally fed adults. *Nutr Clin Pract* 2009;24:261-73.
95. Collard HR, Saint S, Matthay MA. Prevention of ventilator-associated pneumonia: An evidence-based systematic review. *Ann Intern Med* 2003;138:494-501.
96. Drakulovic MB, Torres A, Bauer TT, Nicolas JM, Nogué S, Ferrer M. Supine body position as a risk factor for nosocomial pneumonia in mechanically ventilated patients: A randomised trial. *Lancet* 1999;354(9193):1851-8.
97. van Nieuwenhoven CA, Vandebroucke-Grauls C, van Tiel FH, Joore HC, van Schijndel RJS, van der Tweel I; *et al.* Feasibility and effects of the semirecumbent position to prevent ventilator-associated pneumonia: A randomized study. *Crit Care Med* 2006;34:396-402.
98. Bonten MJ, Gaillard CA, van der Hulst R, De Leeuw PW, van der Geest S, Stobberingh EE, Soeters PB. Intermittent enteral feeding: the influence on respiratory and digestive tract colonization in mechanically ventilated intensive-care-unit patients. *Am J Respir Crit Care Med* 1996;154(2 Pt 1):394-9.
99. Steevens EC, Lipscomb AF, Poole GV, Sacks GS. Comparison of continuous vs intermittent nasogastric enteral feeding in trauma patients: Perceptions and practice. *Nutr Clin Pract* 2002;17:118-22.
100. MacLeod JB, Lefton, J, Houghton D, Roland C, Doherty J, Cohn SM, Barquist ES. Prospective randomized control trial of intermittent versus continuous gastric feeds for critically ill trauma patients. *J Trauma* 2007;63:57-61.

101. Ho KM, Dobb GJ, Webb SA. A comparison of early gastric and post-pyloric feeding in critically ill patients: A meta-analysis. *Intensive Care Med* 2006;32:639-49.
102. Marik PE, Zaloga GP. Gastric versus post-pyloric feeding: A systematic review. *Crit Care* 2003;7:R46-R51.
103. Heyland DK, Drover JW, Dhaliwal R, Greenwood J. Optimizing the benefits and minimizing the risks of enteral nutrition in the critically ill: Role of small bowel feeding. *JPEN J Parenter Enteral Nutr* 2002;26(6 Suppl):S51-S55.
104. Deane AM, Dhaliwal R, Day AG, Ridley EJ, Davies AR, Heyland DK. Comparisons between intragastric and small intestinal delivery of enteral nutrition in the critically ill: A systematic review and meta-analysis. *Crit Care* 2014;17:R125.
105. Alhazzani W, Almasoud A, Jaeschke R, Lo BW, Sindi A, Altayyar S, Fox-Robichaud A. Small bowel feeding and risk of pneumonia in adult critically ill patients: A systematic review and meta-analysis of randomized trials. *Crit Care* 2013;17:R127.
106. McClave SA, Sexton LK, Spain DA, Adams JL, Owens NA, Sullins MB; *et al.* Enteral tube feeding in the intensive care unit: Factors impeding adequate delivery. *Crit Care Med* 1999;27:1252-6.
107. O'Leary-Kelley CM, Puntillo KA, Barr J, Stotts N, Douglas MK. Nutritional adequacy in patients receiving mechanical ventilation who are fed enterally. *Am J Crit Care* 2005;14: 222-31.
108. Montejo JC, for the Nutritional and Metabolic Working Group of the Spanish Society of Intensive Care Medicine and Coronary Units. Enteral nutrition-related gastrointestinal complications in critically ill patients: A multicenter study. *Crit Care Med* 1999;27:1447-53.
109. Parrish C, McClave S. Checking gastric residual volumes: A practice in search of science? *Practical Gastroenterology* 2008;10:33-47.
110. Montejo JC, Minambres E, Bordeje L, Mesejo A, Acosta J, Heras A; *et al.* Gastric residual volume during enteral nutrition in ICU patients: The REGANE study. *Intensive Care Med* 2010; 36:1386-93.
111. Pinilla JC, Samphire J, Arnold C, Liu L, Thiessen, B. Comparison of gastrointestinal tolerance to two enteral feeding protocols in critically ill patients: A prospective, randomized controlled trial. *JPEN J Parenter Enteral Nutr* 2001;25:81-6.
112. McClave SA, Lukan JK, Stefater JA, Lowen CC, Looney SW, Matheson PJ; *et al.* Poor validity of residual volumes as a marker for risk of aspiration in critically ill patients. *Crit Care Med* 2005;33:324-30.
113. Chapman MJ. Gastric emptying and nutrition in the critically ill patient. *Crit Care Resusc* 2003;5:11-3.
114. Dive A, Miesse C, Galanti L, Jamart J, Evrard P, Gonzalez M; *et al.* Effect of erythromycin on gastric motility in mechanically ventilated critically ill patients: A double-blind, randomized, placebo-controlled study. *Crit Care Med* 1995;23:1356-62.
115. Marino LV, Kiratu EM, French N, Nathoo N. To determine the effect of metoclopramide on gastric emptying in severe head injuries: A prospective, randomized, controlled clinical trial. *Br J Neurosurg* 2003;17:24-8.
116. Jooste CA, Mustoe J, Collee G. Metoclopramide improves gastric motility in critically ill patients. *Intensive Care Med* 1999;25:464-8.
117. Reignier J, Bensaid S, Perrin-Gachadoat D, Burdin M, Boiteau R, Tenaillon A. Erythromycin and early enteral nutrition in mechanically ventilated patients. *Crit Care Med* 2002; 30:1237-41.

118. Nursal TZ, Erdogan B, Noyan T, Cekinmez M, Atalay B, Bilgin, N. The effect of metoclopramide on gastric emptying in traumatic brain injury. *J ClinNeurosci* 2007;14:344-8.
119. Yavagal DR, Karnad DR, Oak JL. Metoclopramide for preventing pneumonia in critically ill patients receiving enteral tube feeding: A randomized controlled trial. *Crit Care Med* 2000;28:1408-11.
120. Meissner W, Dohrn B, Reinhart K. Enteral naloxone reduces gastric tube reflux and frequency of pneumonia in critical care patients during opioid analgesia. *Crit Care Med* 2003; 31:776-80.
121. Chapman MJ, Chapman MJ, Fraser RJ, Kluger MT, Buist MD, De Nichilo DJ. Erythromycin improves gastric emptying in critically ill patients intolerant of nasogastric feeding. *Crit Care Med* 2000;28:2334-7.
122. Berne JD, Norwood SH, McAuley CE, Vallina VL, Villareal D, Weston J, McClarty J. Erythromycin reduces delayed gastric emptying in critically ill trauma patients: A randomized, controlled trial. *J Trauma* 2002;53:422-5.
123. MacLaren R, Kiser TH, Fish DN, Wischmeyer PE. Erythromycin vs. metoclopramide for facilitating gastric emptying and tolerance to intragastric nutrition in critically ill patients. *JPEN J Parenter Enteral Nutr* 2008;32:412-9.
124. Nguyen NQ, Chapman M, Fraser RJ, Bryant LK, Burgstad C, Holloway RH. Prokinetic therapy for feed intolerance in critical illness: One drug or two? *Crit Care Med* 2007;35: 2561-7.
125. Nguyen NQ, Chapman MJ, Fraser RJ, Bryant LK, Holloway RH. Erythromycin is more effective than metoclopramide in the treatment of feed intolerance in critical illness. *Crit Care Med* 2007;35:483-9.
126. Preiser JC, Ledoux D. The use of protocols for nutritional support is definitely needed in the intensive care unit. *Crit Care Med* 2004;32:2354-5.
127. Montejo González JC, Estébanez Montiel B. Complicaciones gastrointestinales en el paciente crítico. *Nutrición Hospitalaria [España]* 2007;22:56-62.
128. Jenkins ME, Gottschlich MM, Warden GD. Enteral feeding during operative procedures in thermal injuries. *J Burn Care Rehabil* 1994;15:199-205.
129. Barr J, Hecht M, Flavin KE, Khorana A, Gould MK. Outcomes in critically ill patients before and after the implementation of an evidence-based nutritional management protocol. *Chest* 2004;125:1446-57.
130. Silk DB, Fairclough PD, Clark ML, Hegarty JE, Marrs TC, Addison JM; *et al.* Use of a peptide rather than free amino acid nitrogen source in chemically defined "elemental" diets. *JPEN J Parenter Enteral Nutr* 1997;4:548-53.
131. Ziegler F, Ollivier JM, Cynober L, Masini JP, Coudray-Lucas C, Levy E, Giboudeau J. Efficiency of enteral nitrogen support in surgical patients: Small peptides *vs.* non-degraded proteins. *Gut* 1990;31:1277-83.
132. Wagner DR, Elmore MF, Knoll DM. Evaluation of "closed" *vs.* "open" systems for the delivery of peptide-based enteral diets. *JPEN J Parenter Enteral Nutr* 1994;18:453-7.
133. Mowatt-Larsen CA, Brown RO, Wojtysiak SL, Kudsk KA. Comparison of tolerance and nutritional outcome between a peptide and a standard enteral formula in critically ill, hypoalbuminemic patients. *JPEN J Parenter Enteral Nutr* 1992;16:20-4.
134. Heimbigner DC, Geels WJ, Bilbrey J, Redden DT, Keeney C. Effects of small-peptide and whole-protein enteral feedings on serum proteins and diarrhea in critically ill patients: A randomized trial. *JPEN J Parenter Enteral Nutr* 1997;21:162-7.

135. Brinson RR, Kolts BE. Diarrhea associated with severe hypoalbuminemia: A comparison of a peptide-based chemically defined diet and standard enteral alimentation. *Crit Care Med* 1988;16:130-6.
136. Meredith JW, Ditesheim JA, Zaloga GP. Visceral protein levels in trauma patients are greater with peptide diet than with intact protein diet. *J Trauma* 1990;30:825-8.
137. de Aguilar-Nascimento JE, Prado Silveira BR, Dock-Nascimento DB. Early enteral nutrition with whey protein or casein in elderly patients with acute ischemic stroke: A double-blind randomized trial. *Nutrition* 2011;27:440-4.
138. Heys SD, Walker LG, Smith I, Eremin O. Enteral nutritional supplementation with key nutrients in patients with critical illness and cancer: A meta-analysis of randomized controlled clinical trials. *Ann Surg* 1999;229:467-77.
139. Beale RJ, Bryg DJ, Bihari DJ. Immunonutrition in the critically ill: A systematic review of clinical outcome. *Crit Care Med* 1999;27:2799-805.
140. Heyland DK, Novak F, Drover JW, Jain M, Su X, Suchner U. Should immunonutrition become routine in critically ill patients? A systematic review of the evidence. *JAMA* 2001; 286: 944-53.
141. Montejo JC, Zarazaga A, López-Martínez J, Urrutia G, Roqué M, Blesa AL; *et al.* Immunonutrition in the intensive care unit. A systematic review and consensus statement. *Clin Nutr* 2003;22:221-33.
142. Waitzberg DL, Saito H, Plank LD, Jamieson GG, Jagannath P, Hwang TL. Postsurgical infections are reduced with specialized nutrition support. *World J Surg* 2006;30:1592-604.
143. Marik PE, Zaloga GP. Immunonutrition in critically ill patients: A systematic review and analysis of the literature. *Intensive Care Med* 2008;34:1980-90.
144. Bower RH, Cerra FB, Bershadsky B, Licari JJ, Hoyt DB, Jensen GL; *et al.* Early enteral administration of a formula (Impact) supplemented with arginine, nucleotides, and fish oil in intensive care unit patients: Results of a multicenter, prospective, randomized, clinical trial. *Crit Care Med* 1995;23:436-49.
145. Bertolini G, Iapichino G, Radrizzani D, Facchini R, Simini B, Bruzzone P; *et al.* Early enteral immunonutrition in patients with severe sepsis: Results of an interim analysis of a randomized multicentre clinical trial. *Intensive Care Med* 2003;29:834-40.
146. Lorente JA, Landín L, De Pablo R, Renes E, Liste D. L-arginine pathway in the sepsis syndrome. *Crit Care Med* 1993;21:1287-95.
147. Lorente JA, Landín L, Renes E, De Pablo R, Jorge P, Ródena E, Liste D. Role of nitric oxide in the hemodynamic changes of sepsis. *Crit Care Med* 1993;21:759-67.
148. Schneider H, Atkinson SW. Immunonutrition -Fact or fad. *J Nutr Health Aging* 2000; 4:120-3.
149. Heyland D, Muscedere J, Wischmeyer PE, Cook D, Jones G, Albert M; *et al.* A randomized trial of glutamine and antioxidants in critically ill patients. *N Engl J Med* 2013; 368:1489-97.
150. van Zanten AR, Sztark F, Kaisers UX, Zielmann S, Felbinger TW, Sablotzki AR; *et al.* High-protein enteral nutrition enriched with immune modulating nutrients vs. standard high-protein enteral nutrition and nosocomial infections in the ICU: A randomized clinical trial. *JAMA* 2014;312:514-24.
151. Atkinson S, Sieffert E, Bihari D; for the Guy's Hospital Intensive Care Group. A prospective, randomized, double-blind, controlled clinical trial of enteral immunonutrition in the critically ill. *Crit Care Med* 1998;26:1164-72.

152. Gadek JE, DeMichele SJ, Karlstad MD, Pacht ER, Donahoe M, Albertson TE; *et al*; for the Enteral Nutrition in ARDS Study Group. Effect of enteral feeding with eicosapentaenoic acid, gamma linolenic acid, and antioxidants in patients with acute respiratory distress syndrome. Crit Care Med 1999;27:1409-20.
153. Pontes-Arruda A, Aragao AM, Albuquerque J.D. Effects of enteral feeding with eicosapentaenoic acid, gamma-linolenic acid, and antioxidants in mechanically ventilated patients with severe sepsis and septic shock. Crit Care Med 2006;34:2325-33.
154. Singer P, Theilla M, Fisher H, Gibstein L, Grozovski E, Cohen J. Benefit of an enteral diet enriched with eicosapentaenoic acid and gamma-linolenic acid in ventilated patients with acute lung injury. Crit Care Med 2006;34:1033-8.
155. Rice TW, Wheeler AP, Thompson BT, DeBoisblanc BP, Steingrub J, Rock P; for the NHLBI ARDS Clinical Trials Network. Enteral omega-3 fatty acid, gamma-linolenic acid, and antioxidant supplementation in acute lung injury. JAMA 2011;306:1574-81.
156. Zhu D, Zhang Y, Li S, Gan L, Feng H, Nie W. Enteral omega-3 fatty acid supplementation in adult patients with acute respiratory distress syndrome: A systematic review of randomized controlled trials with meta-analysis and trial sequential analysis. Intensive Care Med 2014;40:504-12.
157. Parry-Billings M, Calder PC, Newsholme EA, Evans J. Does glutamine contribute to immunosuppression after major burns? Lancet 1990;336(8714):523-5.
158. Roth E, Funovics J, Mühlbacher F, Schemper M, Mauritz W, Sporn P, Fritsch, A. Metabolic disorders in severe abdominal sepsis: Glutamine deficiency in skeletal muscle. Clin Nutr 1982;1:25-41.
159. Oudemans-van Straaten HM, Bosman RJ, Treskes M, van der Spoel HJI, Zandstra DF. Plasma glutamine depletion and patient outcome in acute ICU admissions. Intensive Care Med 2001;27:84-90.
160. Cetinbas F, Yelken B, Gulbas Z. Role of glutamine administration on cellular immunity after total parenteral nutrition enriched with glutamine in patients with systemic inflammatory response syndrome. J Crit Care 2010;25:661-6.
161. Kelly D, Wischmeyer PE. Role of L-glutamine in critical illness: New insights. Curr Opin Clin Nutr Metab Care 2003;6:217-22.
162. Singleton KD, Beckey VE, Wischmeyer PE. Glutamine prevents activation of NF-kappa B and stress kinase pathways, attenuates inflammatory cytokine release and prevents acute respiratory distress syndrome (ARDS) following sepsis. Shock 2005;24:583-9.
163. Bakalar B, Duska F, Pachl J, Fric M, Otahal M, Pazout J, Andel M. Parenterally administered dipeptide alanyl-glutamine prevents worsening of insulin sensitivity in multiple-trauma patients. Crit Care Med 2006;34:381-6.
164. Déchelotte P, Hasselmann M, Cynober L, Allaouchiche B, Coëffier M, Hecketswiler B; *et al*. L-alanyl-L-glutamine dipeptide-supplemented total parenteral nutrition reduces infectious complications and glucose intolerance in critically ill patients: The French controlled, randomized, double-blind, multicenter study. Crit Care Med 2006;34:598-604.
165. Zhou YP, Jiang ZM, Sun YH, He GZ, Shu H. The effects of supplemental glutamine dipeptide on gut integrity and clinical outcome after major escharectomy in severe burns: A randomized, double-blind, controlled clinical trial. Clin Nutr Suppl 2004;1:55-60.
166. Griffiths RD, Jones C, Palmer TE. Six-month outcome of critically ill patients given glutamine-supplemented parenteral nutrition. Nutrition 1997;13:295-302.

167. Powell-Tuck J, Jamieson CP, Bettany GEA, Obeid O, Fawcett HV, Archer C, Murphy DL. A double blind, randomised, controlled trial of glutamine supplementation in parenteral nutrition. *Gut* 1999;45:82-8.
168. Fuentes Orozco C, Anaya Prado R, González Ojeda A, Arenas Márquez H, Cabrera Pivaral C, Cervantes Guevara G, Barrera Zepeda LM. L-alanyl-L-glutamine-supplemented parenteral nutrition improves infectious morbidity in secondary peritonitis. *Clin Nutr* 2004; 23:13-21.
169. Xian-Li H, Qing-Jiu M, Jian-Guo L, Yan-Kui C, Xi-Lin D. Effect of total parenteral nutrition (TPN) with and without glutamine dipeptide supplementation on outcome in severe acute pancreatitis (SAP). *Clin Nutr Suppl* 2004; 1:43-7.
170. Estívariz CF, Griffith DP, Luo M, Szeszycki EE, Bazargan N, Dave N; *et al.* Efficacy of parenteral nutrition supplemented with glutamine dipeptide to decrease hospital infections in critically ill surgical patients. *JPEN J Parenter Enteral Nutr* 2008;32:389-402.
171. Novak F, Heyland DK, Avenell A, Drover JW, Su X. Glutamine supplementation in serious illness: A systematic review of the evidence. *Crit Care Med* 2002;30:2022-9.
172. Wischmeyer PE, Dhaliwal R, McCall M, Ziegler TR, Heyland DK. Parenteral glutamine supplementation in critical illness: A systematic review. *Crit Care* 2014;18:R76.
173. Critical Care Nutrition Committee. Canadian Clinical Practice Guidelines. Composition of Parenteral Nutrition: Glutamine. Montreal: 2013. Disponible en: <http://www.criticalcarenutrition.com/docs/cpgs2012/3.3a.pdf>. Fecha de última visita: 11 de Noviembre del 2015.
174. Chen QH, Yang Y, He HL, Xie JF, Cai SX, Liu AR; *et al.* The effect of glutamine therapy on outcomes in critically ill patients: A meta-analysis of randomized controlled trials. *Crit Care* 2014;18:R8.
175. Tao KM, Li XQ, Yang LQ, Yu WF, Lu ZJ, Sun YM, Wu FX. Glutamine supplementation for critically ill adults. *Cochrane Database Syst Rev* 2014;9:CD010050.
176. Garrel D, Patenaude J, Nedelec B, Samson L, Dorais J, Champoux J; *et al.* Decreased mortality and infectious morbidity in adult burn patients given enteral glutamine supplements: A prospective, controlled, randomized clinical trial. *Crit Care Med* 2003;31:2444-9.
177. Zhou YP, Jiang ZM, Sun YH, Wang XR, Ma EL, Wilmore, D. The effect of supplemental enteral glutamine on plasma levels, gut function, and outcome in severe burns: A randomized, double-blind, controlled clinical trial. *JPEN J Parenter Enteral Nutr* 2003;27:241-5.
178. Peng X, Yan H, You Z, Wang P, Wang S. Effects of enteral supplementation with glutamine granules on intestinal mucosal barrier function in severe burned patients. *Burns* 2004;30:135-9.
179. Houdijk AP, Rijnsburger ER, Jansen J, Wesdorp RI, Weiss JK, McCamish MA; *et al.* Randomised trial of glutamine-enriched enteral nutrition on infectious morbidity in patients with multiple trauma. *Lancet* 1998;352(9130):7726.
180. McQuiggan M, Kozar R, Sailors RM, Ahn C, McKinley B, Moore F. Enteral glutamine during active shock resuscitation is safe and enhances tolerance of enteral feeding. *JPEN J Parenter Enteral Nutr* 2008;32:28-35.
181. Conejero R, Bonet A, Grau T, Esteban A, Mesejo A, Montejo JC; *et al.* Effect of a glutamine-enriched enteral diet on intestinal permeability and infectious morbidity at 28 days in critically ill patients with systemic inflammatory response syndrome: A randomized, single-blind, prospective, multicenter study. *Nutrition* 2002;18:716-21.
182. Jones C, Palmer TE, Griffiths RD. Randomized clinical outcome study of critically ill patients given glutamine-supplemented enteral nutrition. *Nutrition* 1999;15:108-15.

183. Hall JC, Dobb G, Hall J, de Sousa R, Brennan L, McCauley R. A prospective randomized trial of enteral glutamine in critical illness. *Intensive Care Med* 2003;29:1710-6.
184. Luo M, Bazargan N, Griffith DP, Estívariz CF, Leader LM, Easley KA; *et al.* Metabolic effects of enteral versus parenteral alanyl-glutamine dipeptide administration in critically ill patients receiving enteral feeding: A pilot study. *Clin Nutr* 2008;27:297-306.
185. Critical Care Nutrition Committee. Canadian Clinical Practice Guidelines. Composition of Enteral Nutrition: Glutamine. Montreal: 2013. Disponible en: <http://www.criticalcarenutrition.com/docs/cpgs2012/3.3a.pdf>. Fecha de última visita: 13 de Diciembre del 2015.
186. Goeters C, Wenn A, Mertes N, Wempe C, van Aken H, Stehle P, Bone HG. Parenteral L-alanyl-L-glutamine improves 6-month outcome in critically ill patients. *Crit Care Med* 2002;30:2032-7.
187. Heller AR, Rössler S, Litz RJ, Stehr SN, Heller SC, Koch R, Koch T. Omega-3 fatty acids improve the diagnosis-related clinical outcome. *Crit Care Med* 2006;34:972-9.
188. Tappy L, Berger MM, Schwarz JM, Schneiter P, Kim S, Revelly P, Chioléro R. Metabolic effects of parenteral nutrition enriched with n-3 polyunsaturated fatty acids in critically ill patients. *Clin Nutr* 2006;25: 588-95.
189. Wichmann MW, Thul P, Czarnetzki HD, Morlion BJ, Kemen M, Jauch KW. Evaluation of clinical safety and beneficial effects of a fish oil containing lipid emulsion (Lipoplus, MLF541): Data from a prospective, randomized, multicenter trial. *Crit Care Med* 2007; 35:700-6.
190. Wang X, Li W, Li N, Li J. Omega-3 fatty acids-supplemented parenteral nutrition decreases hyperinflammatory response and attenuates systemic disease *sequelae* in severe acute pancreatitis: A randomized and controlled study. *JPEN J Parenter Enteral Nutr* 2008;32:236-41.
191. Friescke S, Lotze C, Köhler J, Heinrich A, Felix SB, Abel P. Fish oil supplementation in the parenteral nutrition of critically ill medical patients: A randomised controlled trial. *Intensive Care Med* 2008;34:1411-20.
192. Manzanares W, Dhaliwal R, Jurewitsch B, Stapleton RD, Jeejeebhoy KN, Heyland DK. Parenteral fish oil lipid emulsions in the critically ill. A systematic review and meta-analysis. *JPEN J Parenter Enteral Nutr* 2014;38:20-8.
193. Goodyear-Bruch C, Pierce JD. Oxidative stress in critically ill patients. *Am J Crit Care* 2002;11:543-51.
194. Forceville X, Vitoux D, Gauzit R, Combes A, Lahilaire P, Chappuis P. Selenium, systemic immune response syndrome, sepsis, and outcome in critically ill patients. *Crit Care Med* 1998; 26:1536-44.
195. Berger MM, Shenkin A. Vitamins and trace elements: Practical aspects of supplementation. *Nutrition* 2006;22:952-5.
196. Berger MM. Can oxidative damage be treated nutritionally? *Clin Nutr* 2005;24:172-83.
197. Mishra V, Baines M, Perry SE, McLaughlin PJ, Carson J, Wenstone R, Shenkin A. Effect of selenium supplementation on biochemical markers and outcome in critically ill patients. *Clin Nutr* 2007;26:41-50.
198. Kazda A, Brodská H, Valenta J, Vinglerová M, Hendl J, Stach Z, Slabá I. Selenium and its substitution in critically ill patients. *Critical Care* 2006;10(Suppl 1):P187.
199. Berger M, Reymond M, Shenkin A, Rey F, Wardle C, Cayeux C; *et al.* Influence of selenium supplements on the post-traumatic alterations of the thyroid axis: A placebo-controlled trial. *Intensive Care Med* 2001;27:91-100.

200. Berger MM, Soguel L, Shenkin A, Revelly JP, Pinget C, Baines M, Chioléro RL. Influence of early antioxidant supplements on clinical evolution and organ function in critically ill cardiac surgery, major trauma, and subarachnoid hemorrhage patients. Crit Care 2008;12:R101.
201. Angstwurm MW, Engelmann L, Zimmermann T, Lehmann C, Spes CH, Abel P; et al. Selenium in Intensive Care (SIC): Results of a prospective randomized, placebo-controlled, multiple-center study in patients with severe systemic inflammatory response syndrome, sepsis, and septic shock. Crit Care Med 2007;35:118-26.
202. Forceville X, Laviolle B, Annane D, Vitoux D, Bleichner G, Korach JM; et al. Effects of high doses of selenium, as sodium selenite, in septic shock: A placebo-controlled, randomized, double-blind, phase II study. Crit Care 2007;11:R73.
203. Andrews PJ, Avenell A, Noble DW, Campbell MK, Croal BL, Simpson WG; et al. Randomised trial of glutamine, selenium, or both, to supplement parenteral nutrition for critically ill patients. BMJ 2011;342:D1542.
204. Berger MM, Spertini F, Shenkin A, Wardle C, Wiesner L, Schindler C, Chiolero RL. Trace element supplementation modulates pulmonary infection rates after major burns: A double-blind, placebo-controlled trial. Am J ClinNutr 1998;68:365-71.
205. Nathens AB, Neff MJ, Jurkovich GJ, Klotz P, Farver K, Ruzinski JT; et al. Randomized, prospective trial of antioxidant supplementation in critically ill surgical patients. Ann Surg 2002;236: 814-22.
206. Heyland DK, Dhaliwal R, Suchner U, Berger MM. Antioxidant nutrients: A systematic review of trace elements and vitamins in the critically ill patient. Intensive Care Med 2005; 31:327-37.
207. Martinuzzi ALN, Ferraresi E., Santana Porbén S, Alcántara S, Alonso M. Diarrea en el paciente crítico: Su actualidad. RCAN Rev Cubana Aliment Nutr 2012;22:120-34.
208. Chang SJ, Huang HH. Diarrhea in enterally fed patients: Blame the diet? Curr Opin Clin Nutr Metab Care 2013;16:588-94.
209. Borges SL, Pinheiro BDV, Pace FHDL, Chebli JMF. Nosocomial diarrhea in the intensive care unit: Incidence and risk factors. Arq Gastroenterol 2008;45:117-23.
210. Btaiche IF, Chan LN, Pleva M, Kraft MD. Critical illness, gastrointestinal complications, and medication therapy during enteral feeding in critically ill adult patients. Nutr Clin Pract 2010;25:32-49.
211. Cohen SH, Gerding DN, Johnson S, Kelly CP, Loo VG, McDonald LC; et al. Clinical practice guidelines for *Clostridium difficile* infection in adults: 2010 update by the Society for Healthcare Epidemiology of America (SHEA) and the Infectious Diseases Society of America (IDSA). Infect Control Hosp Epidemiol 2010;31:431-55.
212. Wiesen P, van Goossum A, Preiser JC. Diarrhoea in the critically ill. Curr Opin Crit Care 2006;12:149-54.
213. Kyne L, Warny M, Qamar A, Kelly CP. Asymptomatic carriage of *Clostridium difficile* and serum levels of IgG antibody against toxin A. N Engl J Med 2000;342:390-7.
214. Miggins M, Hasan A, Hohmann S, Southwick F, Casella G, Schain D; et al. The potential influence of common viral infections diagnosed during hospitalization among critically ill patients in the United States. PLoS One 2011;6(4):e18890.
215. Chittawatanarat K, Pokawinpudisnun P, Polbhakdee Y. Mixed fibers diet in surgical ICU septic patients. Asia Pac J Clin Nutr 2010;19:458-64.
216. Santana Porbén S. Los probióticos en la práctica médica: Razones para su uso. RCAN Rev Cubana Aliment Nutr 2009;19:124-32.

217. Frohmader TJ, Chaboyer WP, Robertson IK, Gowardman J. Decrease in frequency of liquid stool in enterally fed critically ill patients given the multispecies probiotic VSL#3: A pilot trial. *Am J Crit Care* 2010;19:e1-e11.
218. Bleichner G, Blehaut H, Mentec H, Moyse D. *Saccharomyces boulardii* prevents diarrhea in critically ill tube-fed patients. A multicenter, randomized, double-blind placebo-controlled trial. *Intensive Care Med* 1997;23:517-23.
219. Kotzampassi K, Giamarellos-Bourboulis EJ, Voudouris A, Kazamias P, Eleftheriadis E. Benefits of a synbiotic formula (Synbiotic 2000Forte©) in critically ill trauma patients: Early results of a randomized controlled trial. *World J Surg* 2006;30:1848-55.
220. Alberda C, Gramlich L, Meddings J, Field C, McCargar L, Kutsogiannis D; *et al.* Effects of probiotic therapy in critically ill patients: A randomized, double-blind, placebo-controlled trial. *Am J Clin Nutr* 2007;85:816-23.
221. Ferrie S, Daley M. *Lactobacillus GG* as treatment for diarrhea during enteral feeding in critical illness randomized controlled trial. *JPEN J Parenter Enteral Nutr* 2011;35:43-9.
222. Knight DJ, Gardiner D, Banks A, Snape SE, Weston VC, Bengmark S, Girling KJ. Effect of synbiotic therapy on the incidence of ventilator associated pneumonia in critically ill patients: A randomised, double-blind, placebo-controlled trial. *Intensive Care Med* 2009; 35:854-61.
223. Morrow LE, Gogineni V, Malesker MA. Probiotics in the intensive care unit. *Nutr Clin Pract* 2012;27:235-41.
224. Besselink MG, Timmerman HM, Buskens E, Nieuwenhuijs VB, Akkermans LM, Gooszen HG. Probiotic prophylaxis in patients with predicted severe acute pancreatitis (PROPATRIA): Design and rationale of a double-blind, placebo-controlled randomised multicenter trial [ISRCTN38327949]. *BMC Surgery* 2004;4(1):1. Disponible en: <http://bmcsurg.biomedcentral.com/articles/10.1186/1471-2482-4-12>. Fecha de última visita: 12 de Enero del 2106.
225. Morrow LE, Gogineni V, Malesker MA. Synbiotics and probiotics in the critically ill after the PROPATRIA trial. *Curr Opin Clin Nutr Metab Care* 2012;15:147-50.
226. Uchino S, Kellum JA, Bellomo R, Doig GS, Morimatsu H, Morgera S; *et al.* Acute renal failure in critically ill patients: A multinational, multicenter study. *JAMA* 2005;294:813-8.
227. Fiaccadori E, Maggiore U, Rotelli C, Giacosa R, Picetti E, Parenti E; *et al.* Effects of different energy intakes on nitrogen balance in patients with acute renal failure: A pilot study. *Nephrol Dial Transplant* 2005;20:1976-80.
228. Bagshaw SM, Cruz DN, Gibney RN, Ronco C. A proposed algorithm for initiation of renal replacement therapy in adult critically ill patients. *Crit Care* 2009;13:317.
229. Druml W, Bürger U, Kleinberger G, Lenz K, Laggner A. Elimination of amino acids in acute renal failure. *Nephron* 1986;42:62-7.
230. Brochard L, Abroug F, Brenner M, Broccard AF, Danner RL, Ferrer M; *et al.* An official ATS/ERS/ESICM/SCCM/SRLF statement: Prevention and management of acute renal failure in the ICU Patient: An international consensus conference in intensive care medicine. *Am J Respir Crit Care Med* 2010;181:1128-55.
231. Fiaccadori E, Parenti E, Maggiore U. Nutritional support in acute kidney injury. *J Nephrol* 2008;21:645-56.
232. Frankenfield DC, Badellino MM, Reynolds HN, Wiles CE, Siegel JH, Goodarzi S. Amino acid loss and plasma concentration during continuous hemodiafiltration. *JPEN J Parenter Enteral Nutr* 1993;17:551-61.

233. Klein CJ, Moser-Veillon PB, Schweitzer A, Douglass LW, Reynolds HN, Patterson KY, Veillon C. Magnesium, calcium, zinc, and nitrogen loss in trauma patients during continuous renal replacement therapy. *J PEN J Parenter Enteral Nutr* 2002;26:77-92.
234. Druml W, Kierdorf HP; for the Working Group for Developing the Guidelines for parenteral nutrition of the German Association for Nutritional Medicine. Parenteral nutrition in patients with renal failure. En: Guidelines on parenteral nutrition. Chapter 17. Ger Med Sci 2009;7. Disponible en: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2795369/>. Fecha de última visita: 13 de Enero del 2016.
235. Bellomo R, Tan HK, Bhonagiri S, Gopal I, Seacombe J, Daskalakis M, Boyce N. High protein intake during continuous hemodiafiltration: Impact on amino acids and nitrogen balance. *Int J Artif Organs* 2002;25:261-8.
236. Scheinkestel CD, Kar L, Marshall K, Bailey M, Davies A, Nyulasi I, Tuxen DV. Prospective randomized trial to assess caloric and protein needs of critically ill, anuric, ventilated patients requiring continuous renal replacement therapy. *Nutrition* 2003;19:909-16.
237. Scheinkestel CD, Adams F, Mahony L, Bailey M, Davies AR, Nyulasi I, Tuxen DV. Impact of increasing parenteral protein loads on amino acid levels and balance in critically ill anuric patients on continuous renal replacement therapy. *Nutrition* 2003;19:733-40.
238. Mesejo A, Acosta JA, Ortega C, Vila J, Fernández M, Ferreres J; *et al.* Comparison of a high-protein disease-specific enteral formula with a high-protein enteral formula in hyperglycemic critically ill patients. *Clin Nutr* 2003;22:295-305.
239. van den Berg B, Bogaard JM, Hop WC. High fat, low carbohydrate, enteral feeding in patients weaning from the ventilator. *Intensive Care Med* 1994;20:470-5.
240. Al-Saady NM, Blackmore CM, Bennett ED. High fat, low carbohydrate, enteral feeding lowers PaCO₂ and reduces the period of ventilation in artificially ventilated patients. *Intensive Care Med* 1989;15:290-5.
241. Alish CJ, Garvey WT, Maki KC, Sacks GS, Hustead DS, Hegazi RA, Mustad VA. A diabetes-specific enteral formula improves glycemic variability in patients with type 2 diabetes. *Diabetes Technol Ther* 2010;12:419-25.
242. Schwingshackl L, Hoffmann G. Long-term effects of low glycemic index/load vs. high glycemic index/load diets on parameters of obesity and obesity-associated risks: A systematic review and meta-analysis. *Nutr Metab Cardiovasc Dis* 2013;23:699-706.